Author Index

Adachi, K., see Suzuki, N. 181
Alley, E. G., see Lynn, B. C. 211
Amato, A., see Gagliardi, L. 409
Ando, S.
——, Waki, H. and Kon, K.
High-performance liquid chromatography of underivatized gangliosides 285
Asotra, S.
——, Mladenov, P. V. and Burke, R. D.
Improved method for benzoyl chloride derivatization of polyamines for high-performance liquid chromatography 227
Aubel, M. T.
—— and Rogers, L. B.
Effects of pretreatment on the enantioselectivity of silica-bound proteins used as high-performance liquid chromatographic stationary phases 99
Aue, W. A., see Siu, K. W. M. 53
——, see Tang, Y.-Z. 69
Bagchi, S. P., see Jindal, S. P. 356
Bakos, J., see Lecomte, L. 416
Baláspiri, L., see Péter, A. 43
Barbieri, L.
——, Stoppa, C. and Bolognesi, A.
Large scale chromatographic purification of ribosome-inactivating proteins 235
Bardalaye, P. C., see Toth, J. P. 335
Basili, A., see Gagliardi, L. 409
Berman, S. S., see Siu, K. W. M. 53
Bolognesi, A., see Barbieri, L. 235
Borner, S.
—— and Brenneisen, R.
Determination of tryptamine derivatives in hallucinogenic mushrooms using high-performance liquid chromatography with photodiode array detection 402
Bossinger, P., see Räder, K. 341
Bowman, P. B., see Clark, C. P. 275
Bramer, A. M., see Hughes, D. E. 296
Brenneisen, R., see Borner, S. 402
Brieva, A., see Rubio, N. 420
Burger, K., see Péter, A. 43
Burke, R. D., see Asotra, S. 227
Cavazzutti, G., see Gagliardi, L. 409
Cheng, W.
—— and Hollis, D.
Flow-rate effect on elution volume in size-exclusion chromatography 9
Chouliotis, A., see Noël, D. 129
Clark, C. P.
——, Snider, B. G. and Bowman, P. B.
High-performance liquid chromatographic method for determining the enantiomeric purity of a benzindene prostaglandin by a diastereomeric separation 275
Clarke, M. L., see Ivin, P. C. 393
Cohen, A. I., see Jemal, M. 360
——, see Wang-Iverson, D. 365
Cohen, I. C., see Lumley, I. D. 115
Cole, K. C., see Noël, D. 129
Costa, M., see Murphy, R. 388
Cutié, S. S.
——, Kallos, G. J. and Smith, P. B.
Separation and characterization of acrylic acid oligomers by nuclear magnetic resonance spectroscopy and thermospray ion-exchange liquid chromatography–mass spectrometry 349
Den, T. G.
—— and Lee, S. L.
Stationary phases. XV. High-performance liquid chromatographic behaviour of quinones on a safrole stationary phase 323
Dix, K. D.
—— and Fritz, J. S.
Simple steam distillation for sample preparation prior to gas chromatographic determination of organic compounds 201
Dyr, J. E.
—— and Suttnar, J.
On the increased retention volume of human haemoglobin in high-performance gel filtration 303
Fanali, S.
——, Sinibaldi, M. and Quaglia, M. G.
Isotachophoretic analysis of amoxycillin and carboxymethylcysteine in pharmaceuticals 441
Fritz, J. S., see Dix, K. D. 201
Furness, J. B., see Murphy, R. 388
Gagliardi, L.
——, Amato, A., Basili, A., Cavazzutti, G. and Tonelli, D.
Determination of sun-screen agents in cosmetic products by reversed-phase high-performance liquid chromatography 409
Gathercole, P. S.
—— and Thiel, P. G.
Liquid chromatographic determination of the cyanoginosins, toxins produced by the cyanobacterium *Microcystis aeruginosa* 435

Geyer, R., see McKerrell, P. J. 399
Gholson, A. R.
——, St. Louis, R. and Hill, H. H.
Determination of silicon number in silylated organic compounds using the hydrogen atmosphere flame ionziation detector 329
Gilpin, R. K., see Yang, S. S. 93
Glatz, Z., see Škabrahová, Z. 372
Hansen, B., see Welinder, B. S. 191
Hatano, H., see Yamamoto, F. M. 21
Hechler, J.-J., see Noël, D. 129
Heil, B., see Lecomte, L. 416
Hill, H. H., see Gholson, A. R. 329
Hirokawa, T.
——, Tsuyoshi, T. and Kiso, Y.
Isotachophoretic determination of mobility and pK_a by means of computer simulation. VI. Mobility and pK_a of sixty anions in methanol 27
Hollis, D., see Cheng, W. 9
Hughes, D. E.
—— and Bramer, A. M.
Anomalous liquid chromatographic results originating from the presence of soap residues and the use of nylon filters 296
Ivin, P. C.
—— and Clarke, M. L.
Isolation of kestoses and nystose from enzyme digests by high-performance liquid chromatography 393
James, L. B.
Amino acid analysis: a fall-off in performance 291
Jemal, M.
—— and Cohen, A. I.
Simultaneous determination of the acid chloride S-benzoyl-3-mercapto-2-methylpropanoyl chloride and residual carboxylic acid by capillary gas chromatography 360
——, see Wang-Iverson, D. 365
Jindal, S. P.
——, Lutz, T. and Bagchi, S. P.
Gas–liquid chromatographic–mass spectrometric determination of 1-methyl-4-phenyl-4-propionoxypiperidine using a stable isotope-labeled analogue as an internal standard 356
Jönsson, J. Å.
—— and Lövkvist, P.
Determination of adsorption isotherms from chromatographic measurements, using the peak maxima method 1
Kahle, V., see Škabrahová, Z. 372
Kallos, G. J., see Cutié, S. S. 349
Kayano, M., see Miwa, T. 316
Kiso, Y., see Hirokawa, T. 27
Kolarz, B. N.
——, Łuczyński, J., Trochimczuk, A. and Wojaczyńska, M.
Gas chromatographic properties of porous polyacrylonitrile and divinylbenzene copolymers 308
Kon, K., see Ando, S. 285
Kondoh, Y.
—— and Takano, S.
Determination of zinc pyrithione in cosmetic products by high-performance liquid chromatography with pre-labelling 255
Kovář, J., see Škabrahová, Z. 372
Kumanotani, J., see Yamauchi, Y. 171
Lecomte, L.
——, Triolet, J., Sinou, D., Bakos, J. and Heil, B.
High-performance liquid chromatographic separation of sulphonated phosphines 416
Lee, S. L., see Den, T. G. 323
Lehrfeld, J.
Simultaneous gas–liquid chromatographic determination of aldoses and alduronic acids 245
Lewis, S., see Wilson, I. D. 445
Liang, W., see Zeng, W. 426
Lövkvist, P., see Jönsson, J. Å. 1
Łuczyński, J., see Kolarz, B. N. 308
Lumley, I. D.
——, Patel, I. and Cohen, I. C.
Evaluation of laboratory-prepared high-performance size-exclusion columns. Applications to food analysis 115
Lutz, T., see Jindal, S. P. 356
Lynn, B. C.
—— and Alley, E. G.
Derivatization of diaminocyclohexanes for determination of *cis* and *trans* isomers by gas chromatography 211
McKerrell, P. J.
——, Saunders, G. A. and Geyer, R.
Determination of 4,4′-methylenebis(2-chloroaniline) in urine by high-performance liquid chromatography 399
Marko-Varga, G.
High-performance liquid chromatographic separation of some mono- and disaccharides with detection by a post-column enzyme reactor and a chemically modified electrode 157
Martin, B. M., see Mussar, K. J. 378
Millot, M.-C.
—— and Sebille, B.
Rapid preparation of bovine mercaptalbumin by means of covalent chromatography on silica-based materials 263

Miwa, T.
——, Miyakawa, T., Kayano, M. and Miyake, Y.
Application of an ovomucoid-conjugated column for the optical resolution of some pharmaceutically important compounds 316
Miyakawa, T., see Miwa, T. 316
Miyake, Y., see Miwa, T. 316
Mladenov, P. V., see Asotra, S. 227
Mücke, H.-W., see Räder, K. 341
Murphy, R.
——, Furness, J. B. and Costa, M.
High-sensitivity phenylthiohydantoin amino acid analysis on-line to a gas phase protein sequencer 388
Murray, G. J., see Mussar, K. J. 378
Mussar, K. J.
——, Murray, G. J., Martin, B. M. and Viswanatha, T.
Rapid chromatographic assay procedure for peptide–N-glycosidase activity 378
Nishikata, M.
Separation of β-amino alcohols as their dansyl derivatives by thin-layer chromatography on a polyamide layer sheet 449
Noël, D.
——, Cole, K. C., Hechler, J.-J., Chouliotis, A. and Overbury, K. C.
High-performance size-exclusion chromatography of fiber-reinforced organic-matrix composites 129
Oshima, R., see Yamauchi, Y. 171
Otagawa, T.
——, Zaromb, S. and Stetter, J. R.
Cryogenic sampling for aniline vapor with subsequent liquid chromatography–electrochemical detection analysis 81
Overbury, K. C., see Noël, D. 129
Patel, I., see Lumley, I. D. 115
Péter, A.
——, Szepesi, G., Baláspiri, L. and Burger, K.
Coordinative interactions in the separation of insulin and its derivatives by high-performance liquid chromatography 43
Quaglia, M. G., see Fanali, S. 441
Räder, K.
——, Wildfeuer, A., Wintersberger, F., Bossinger, P. and Mücke, H.-W.
Characterization of bee venom and its main components by high-performance liquid chromatography 341
Řezanka, T., see Víden, I. 145
Rogers, L. B., see Aubel, M. T. 99
Rokushika, S., see Yamamoto, F. M. 21
Rubio, N.
—— and Brieva, A.
Purification of allergens by high-performance liquid chromatography. VII. Purification of the major allergen of birch pollen (*Betula verrucosa*) 420
Saitoh, K., see Suzuki, N. 181
Saunders, G. A., see McKerrell, P. J. 399
Schrader, P. A.
—— and Sisco, W. R.
High-performance liquid chromatographic analysis of fenretinide in rodent feed 430
Sebille, B., see Millot, M.-C. 263
Senō, M., see Yamauchi, Y. 171
Sharma, R., see Subbaramaia, K. 385
Sinibaldi, M., see Fanali, S. 441
Sinou, D., see Lecomte, L. 416
Sisco, W. R., see Schrader, P. A. 430
Siu, K. W. M.
——, Berman, S. S. and Aue, W. A.
Correlations and constants in unipolar and bipolar electron-capture detection 53
Škabrahová, Z.
——, Kovář, J., Glatz, Z., Turánek, J. and Kahle, V.
High-performance liquid chromatographic purification of the H4 and MH3 isoenzymes of lactate dehydrogenase from beef heart 372
Smith, P. B., see Cutié, S. S. 349
Snider, B. G., see Clark, C. P. 275
Sørensen, H. H., see Welinder, B. S. 191
Stetter, J. R., see Otagawa, T. 81
St. Louis, R., see Gholson, A. R. 329
Stoppa, C., see Barbieri, L. 235
Subbaramaiah, K.
—— and Sharma, R.
Immunoaffinity purification of β-amylase from mustard (*Sinapis alba* L.) cotyledons 385
Suttnar, J., see Dyr, J. E. 303
Suzuki, N.
——, Saitoh, K. and Adachi, K.
Reversed-phase high-performance thin-layer chromatography and column liquid chromatography of chlorophylls and their derivatives 181
Szepesi, G., see Péter, A. 43
Takada, M., see Yamauchi, Y. 171
Takano, S., see Kondoh, Yu. 255
Tang, Y.-Z.
—— and Aue, W. A.
Spectra and response linearization of chlorine and bromine compounds in a copper-sensitized flame photometric detector 69

Thiel, P. G., see Gathercole, P. S. 435
Tonelli, D., see Gagliardi, L. 409
Toth, J. P.
—— and Bardalaye, P. C.
Capillary gas chromatographic separation and mass spectrometric detection of cyromazine and its metabolite melamine 335
Triolet, J., see Lecomte, L. 416
Trochimczuk, A., see Kolarz, B. N. 308.
Tsuyoshi, T., see Hirokawa, T. 27
Tu, G., see Zeng, W. 426
Turánek, J., see Škabrahová, Z. 372
Víden, I.
—— and Řezanka, T.
Capillary gas chromatography–mass spectrometry of aliphatic saturated α,ω-dicarboxylic acid dimethyl esters and direct inlet mass spectrometry of the corresponding free acids 145
Viswanatha, T., see Mussar, K. J. 378
Waki, H., see Ando, S. 285
Wang-Iverson, D.
——, Jemal, M. and Cohen, A. I.
Determination of enantiomeric purity of 3-mercapto-2-methylpropionic acid 365
Welinder, B. S.
——, Sørensen, H. H. and Hansen, B.
Recovery of polypeptides after reversed-phase high-performance liquid chromatography 191
Wildfeuer, A., see Räder, K. 341
Willis, D. E.
Automated pre-column derivatization of amino acids with *o*-phthalaldehyde by a reagent sandwiching technique 217
Wilson, I. D.
—— and Lewis, S.
Separation of ecdysteroids by high-performance thin-layer chromatography using automated multiple development 445
Wintersberger, F., see Räder, K. 341
Wojaczyńska, M., see Kolarz, B. N. 308
Yamamoto, F. M.
——, Rokushika, S. and Hatano, H.
Prediction of thermodynamic retention behaviour in reversed-phase liquid chromatography with aqueous eluents and various concentrations of methanol by using the solubility concept model 21
Yamauchi, Y.
——, Takada, M., Oshima, R., Kumanotani, J. and Senō, M.
Liquid chromatographic analysis of products of reaction between methyl linoleate and 4-*tert.*-butyl-*o*-benzoquinone 171
Yang, S. S.
—— and Gilpin, R. K.
Reordering/resolvation of silica immobilized hydrogen bonding ligands used as stationary phases in high-performance liquid chromatography 93
Zaromb, S., see Otagawa, T. 81
Zeng, W.
——, Liang, W. and Tu, G.
Assay of five corynoline-type alkaloids in *Corydalis bungeana* by reversed-phase high-performance liquid chromatography 426